

ON FIELDS OF DEFINITION OF COMPONENTS OF THE SIEGEL SUPERSINGULAR LOCUS

CHIA-FU YU

ABSTRACT. Recently Ibukiyama proves an explicit formula for the number of certain non-principal polarizations on a superspecial abelian surface, extending his earlier work with Katsura for principal polarizations [Compos. Math. 1994]. As a consequence of Ibukiyama's formula, there exists a geometrically irreducible component of the Siegel supersingular locus which is defined over the prime finite field. In this note we give a direct proof of this result.

1. INTRODUCTION

Let p be a rational prime number, and let $n \geq 1$ be a positive integer. Let $\mathcal{A}_n$ denote the coarse moduli space over $\mathbb{F}_p$ of n -dimensional principally polarized abelian varieties. The supersingular locus of $\mathcal{A}_n \otimes \overline{\mathbb{F}}_p$ is denoted by $\overline{\mathcal{S}}_n$, which is the closed reduced $\overline{\mathbb{F}}_p$ -subscheme consisting of all supersingular points in $\mathcal{A}_n(\overline{\mathbb{F}}_p)$. An abelian variety over $\overline{\mathbb{F}}_p$ is *supersingular* (resp. *superspecial*) if it is isogenous (resp. isomorphic) to a product of supersingular elliptic curves. It is known that $\overline{\mathcal{S}}_n$ is defined over $\mathbb{F}_p$, i.e. the action of the Galois group $\Gamma := \text{Gal}(\overline{\mathbb{F}}_p/\mathbb{F}_p)$ leaves the set $\overline{\mathcal{S}}_n$ stable. The unique model of $\overline{\mathcal{S}}_n$ over $\mathbb{F}_p$ in $\mathcal{A}_n$ is denoted by $\mathcal{S}_n$, which is called the supersingular locus of $\mathcal{A}_n$. The set of irreducible components of $\overline{\mathcal{S}}_n$ is denoted by $\Pi_0(\overline{\mathcal{S}}_n)$, on which Γ operates. An irreducible component $V \in \Pi_0(\overline{\mathcal{S}}_n)$ is defined over $\mathbb{F}_p$ if and only if V is stable under the action of Γ . In this note we give a proof of the following result.

Theorem 1.1 (Li-Oort, Ibukiyama). *There exists an irreducible component of $\overline{\mathcal{S}}_n$ that is defined over $\mathbb{F}_p$.*

In Section 2, we shall give some background knowledge on $\Pi_0(\overline{\mathcal{S}}_n)$ due to Li and Oort [5]. Based on loc. cit., Theorem 1.1 is reduced to non-emptiness of the set of certain polarized superspecial abelian varieties that admit a model defined over $\mathbb{F}_p$ (the set $\Lambda_n(\mathbb{F}_p)$ or $\Sigma_n(\mathbb{F}_p)$ in Subsection 2.1); see Theorem 2.3. It follows that Theorem 1.1 is trivial when n is odd. When n is even, Theorem 1.1 then follows from the following result.

Theorem 1.2. *There exists a polarized superspecial abelian surface (A, λ) over $\mathbb{F}_p$ such that $\ker \lambda = A[F]$, where $F : A \rightarrow A^{(p)} = A$ is the relative Frobenius morphism on A .*

In [1, 2] Ibukiyama gives an explicit formula for the cardinality of $\Sigma_2(\mathbb{F}_p)$, and as a byproduct he obtains Theorem 1.2. The result of Ibukiyama confirms the

Date: August 12, 2016.

2010 Mathematics Subject Classification. 11R52, 11G10.

Key words and phrases. superspecial abelian surface, supersingular locus.

existence of polarized abelian surfaces as in Theorem 1.2. We construct such an polarized abelian surface directly in Section 3. This proves Theorem 1.1 by a different method.

2. PRELIMINARIES AND BACKGROUND

2.1. Finite sets Λ_n and Σ_n . Let $\Gamma := \text{Gal}(\overline{\mathbb{F}}_p/\mathbb{F}_p)$ denote the Galois group over $\mathbb{F}_p$, and put $\Gamma_2 := \text{Gal}(\overline{\mathbb{F}}_p/\mathbb{F}_{p^2})$. The Frobenius automorphism $a \mapsto a^p$ in Γ is denoted by σ_p . For any positive integer $n \geq 1$, let Λ_n denote the set of isomorphism classes of superspecial principally polarized abelian varieties of dimension n over $\overline{\mathbb{F}}_p$. This is a finite set on which Γ acts. It is known that the subgroup Γ_2 acts trivially on Λ_n so the action factors through the quotient $\text{Gal}(\mathbb{F}_{p^2}/\mathbb{F}_p)$. Put

$$\Lambda_n(\mathbb{F}_p) := \Lambda_n^\Gamma \subset \Lambda_n,$$

the subset of elements fixed by σ_p . Let E_0 be a supersingular elliptic curve over $\mathbb{F}_p$ such that the endomorphism π_{E_0} of E_0 satisfies $\pi_{E_0}^2 = -p$. Put $A_0 = E_0^n$ and take λ_0 to be the product of copies of the canonical principal polarization of E_0 . Clearly, (A_0, λ_0) is a superspecial principally polarized abelian variety of dimension n over $\mathbb{F}_p$. In particular, the set $\Lambda_n(\mathbb{F}_p)$ is nonempty.

For any even positive integer $n = 2c$, denote by Σ_n the set of isomorphism classes of superspecial polarized abelian varieties (A, λ) of dimension n over $\overline{\mathbb{F}}_p$ such that $\ker \lambda = A[F]$, where $A^{(p)}$ is the base change of $\text{Spec } \overline{\mathbb{F}}_p \rightarrow \text{Spec } \overline{\mathbb{F}}_p$ induced by σ_p , and $F : A \rightarrow A^{(p)}$ is the relative Frobenius morphism. Clearly, for any $(A, \lambda) \in \Sigma_n$, one has $\deg \lambda = p^n$. The Galois group Γ acts on Σ_n which factors through $\text{Gal}(\mathbb{F}_{p^2}/\mathbb{F}_p)$. Indeed, any superspecial abelian variety over $\overline{\mathbb{F}}_p$ admits a model A'_0 over $\mathbb{F}_{p^2}$ such that any endomorphism of A is defined over $\mathbb{F}_{p^2}$ (for the $\mathbb{F}_{p^2}$ -structure of A'_0), in particular, any polarization of A is defined over $\mathbb{F}_{p^2}$. Thus, the group Γ_2 acts trivially on Σ_n . Similarly, we define

$$\Sigma_n(\mathbb{F}_p) := \Sigma_n^\Gamma \subset \Sigma_n,$$

the subset of elements fixed by σ_p . Unlike Λ_n , it is not clear whether $\Sigma_n(\mathbb{F}_p)$ is nonempty a priori.

Recall that the field of moduli of a polarized abelian variety (A, λ) over $\overline{\mathbb{F}}_p$ is the field of definition of the isomorphism class $[(A, \lambda)]$, that is, the smallest finite field $\mathbb{F}_{p^a}$ such that $(A, \lambda) \simeq (\sigma A, \sigma \lambda)$ over $\overline{\mathbb{F}}_p$ for any $\sigma \in \text{Gal}(\overline{\mathbb{F}}_p/\mathbb{F}_{p^a})$.

Lemma 2.1. *Let (A, λ) be any polarized abelian variety over $\overline{\mathbb{F}}_p$, and suppose that the field of moduli of (A, λ) is $\mathbb{F}_{p^a}$. Then (A, λ) has a model defined over $\mathbb{F}_{p^a}$.*

PROOF. The lemma is proved for A being superspecial [2, Lemma 4.1]. The same proof works for an arbitrary abelian variety; also see [7, Prop. 6.3] for a similar proof. ■

Lemma 2.2. *If an element (A, λ) in Σ_n has a model (A', λ') defined over $\mathbb{F}_p$, then $\ker \lambda' = A'[F]$.*

PROOF. We use the following basic fact: if $H_1 < H_2$ are two finite group schemes over a field and if $\text{rank } H_1 = \text{rank } H_2$, then $H_1 = H_2$. It follows that if $\varphi : A \rightarrow B$ is an isogeny of abelian varieties over k , K/k is a field extension and $\varphi_K : A_K \rightarrow$

B_K is the base change morphism, then $\ker \varphi_K = \ker \varphi \otimes_k K$. Applying this for $F : A \rightarrow A^{(p)}$ and $K/k = \overline{\mathbb{F}_p}/\mathbb{F}_p$, we get

$$\ker \lambda' \otimes \overline{\mathbb{F}_p} = \ker \lambda = A[F] = A'[F] \otimes \overline{\mathbb{F}_p}.$$

By the faithfully flat descent, we have $\ker \lambda' = A'[F]$. ■

Let $\tilde{\Sigma}_n(\mathbb{F}_p)$ be the set of isomorphism classes of superspecial polarized abelian varieties (A, λ) of dimension n (n being even) over $\mathbb{F}_p$ such that $\ker(\lambda) = A[F]$. It follows from Lemmas 2.1 and 2.2 that the natural map $\tilde{\Sigma}_n(\mathbb{F}_p) \rightarrow \Sigma_n(\mathbb{F}_p)$ is surjective. Particularly, the set $\Sigma_n(\mathbb{F}_p)$ is nonempty if and only if so is $\tilde{\Sigma}_n(\mathbb{F}_p)$.

2.2. Fields of definition for components. As in Section 1, we let $\mathcal{A}_n$ denote the coarse moduli space over $\mathbb{F}_p$ of n -dimensional principally polarized abelian varieties, and let $\mathcal{S}_n \subset \mathcal{A}_g$ be the supersingular locus of $\mathcal{A}_g$. Li-Oort proved [5, 4.9, p. 26] that there is a one-to-one correspondence between the set $\Pi_0(\overline{\mathcal{S}}_n)$ of irreducible components of $\overline{\mathcal{S}}_n := \mathcal{S}_n \otimes \overline{\mathbb{F}_p}$ and either the set Λ_n or the set Σ_n according as n is odd or even. The Galois group Γ operates on $\Pi_0(\overline{\mathcal{S}}_n)$ as well as on the sets Λ_n and Σ_n . We claim that the correspondence in loc. cit. respects the action of Γ . Let $V \in \Pi_0(\overline{\mathcal{S}}_n)$ be an irreducible component. It is proved in [5, 4.9 (iii), p. 26] that the subset U of supergeneral abelian varieties in the sense of Li and Oort (i.e. with $a = 1$) is an open and dense subset in V . For any point (B, λ_B) in U , there is a unique “PFTQ” (polarized flag type quotient) up to equivalence [5, Chapter 3]

$$(A_{n-1}, \eta_{n-1}) \rightarrow (A_{n-2}, \eta_{n-2}) \rightarrow \cdots \rightarrow (A_0, \eta_0)$$

such that $(A_0, \eta_0) \simeq (B, \lambda_B)$ and $(A_{n-1}, \eta_{n-1}) \simeq (A, p^m \lambda)$ for a superspecial polarized abelian variety (A, λ) that lies in Λ_n if n is odd and in Σ_n if n is even. The pair (A, λ) depends only on U up to isomorphism, and corresponds to the component V . It is then clear from this construction that for any $\sigma \in \Gamma$ the conjugate ${}^\sigma U$ of U corresponds to the isomorphism class $[({}^\sigma A, {}^\sigma \lambda)]$. This proves the claim. Note that the isomorphism class $[({}^\sigma A, {}^\sigma \lambda)]$ is defined over $\mathbb{F}_p$ if and only if (A, λ) admits a model defined over $\mathbb{F}_p$ (Lemma 2.1). Thus, we have the following consequence of results of Li and Oort.

Theorem 2.3 (Li-Oort, cf. [2, Theorem 4.4]).

- (1) Every irreducible component $V \subset \overline{\mathcal{S}}_n$ is defined over $\mathbb{F}_{p^2}$.
- (2) Let $\Pi_0(\overline{\mathcal{S}}_n)(\mathbb{F}_p)$ be the set of irreducible components of $\overline{\mathcal{S}}_n$ that are defined over $\mathbb{F}_p$. Then

$$(2.1) \quad \Pi_0(\overline{\mathcal{S}}_n)(\mathbb{F}_p) \simeq \begin{cases} \Lambda_n(\mathbb{F}_p) & \text{if } n \text{ is odd;} \\ \Sigma_n(\mathbb{F}_p) & \text{if } n \text{ is even.} \end{cases}$$

2.3. Theorem 1.2 implies Theorem 1.1. It follows from Theorem 2.3 that Theorem 1.1 is equivalent to that $\Lambda_n(\mathbb{F}_p)$ is nonempty if n is odd and $\Sigma_n(\mathbb{F}_p)$ is nonempty if n is even. Non-emptiness of $\Lambda_n(\mathbb{F}_p)$ is obvious. Thus, it reduces to prove non-emptiness of $\Sigma_n(\mathbb{F}_p)$. But this following from non-emptiness of $\Sigma_2(\mathbb{F}_p)$, which is equivalent to Theorem 1.2 by Lemma 2.1.

2.4. Explicit formula for $|\Sigma_2(\mathbb{F}_p)|$. Let χ be the quadratic character associated to the real quadratic field $\mathbb{Q}(\sqrt{p})/\mathbb{Q}$, and let $B_{2,\chi}$ be the second generalized Bernoulli number. For any algebraic number α , we write $h(\alpha)$ for the class number of the number field $\mathbb{Q}(\alpha)$.

Theorem 2.4 (Ibukiyama [1, Theorem 5.2]). *One has*

$$|\Sigma_2(\mathbb{F}_p)| = 1, 1, 1, \text{ for } p = 2, 3, 5, \text{ respectively.}$$

For $p \geq 7$, if $p \equiv 1 \pmod{4}$, then

$$(2.2) \quad |\Sigma_2(\mathbb{F}_p)| = \frac{1}{2^5 \cdot 3} \left(9 - 2 \left(\frac{2}{p} \right) \right) B_{2,\chi} + \frac{1}{2^4} h(\sqrt{-p}) \\ + \frac{1}{2^3} h(\sqrt{-2p}) + \frac{1}{2^2 \cdot 3} \left(3 + \left(\frac{2}{p} \right) \right) h(\sqrt{-3p});$$

if $p \equiv 3 \pmod{4}$, then

$$(2.3) \quad |\Sigma_2(\mathbb{F}_p)| = \frac{1}{2^5 \cdot 3} B_{2,\chi} + \frac{1}{2^4} \left(1 - \left(\frac{2}{p} \right) \right) h(\sqrt{-p}) \\ + \frac{1}{2^3} h(\sqrt{-2p}) + \frac{1}{2^2 \cdot 3} h(\sqrt{-3p}).$$

Non-emptiness of $\Sigma_2(\mathbb{F}_p)$ (or Theorem 1.2) follows immediately from Theorem 2.4, and hence Theorem 1.1 follows.

3. CONSTRUCTION OF CERTAIN SUPERSPECIAL POLARIZED ABELIAN VARIETIES

In this section we include some results concerning non-emptiness of the set $\tilde{\Sigma}_n(\mathbb{F}_p)$. The Frobenius endomorphism of an abelian variety A over $\mathbb{F}_q$ will be denoted by π_A . Let $\tilde{\Sigma}'_n(\mathbb{F}_p)$ be the set of isomorphism classes of superspecial abelian varieties (A, λ) of dimension n over $\mathbb{F}_p$ such that $\pi_A^2 = -p$ and $\ker \lambda = A[F]$. It is clear that non-emptiness of $\tilde{\Sigma}'_n(\mathbb{F}_p)$ implies that of $\tilde{\Sigma}_n(\mathbb{F}_p)$.

Lemma 3.1. *Assume that the integer $n = 2c$ is divisible by 4. Then there is a superspecial polarized abelian variety (A', λ') in $\tilde{\Sigma}'_n(\mathbb{F}_p)$.*

PROOF. Choose a superspecial abelian variety A_1 over $\mathbb{F}_{p^2}$ of dimension c such that $\pi_{A_1} = -p$. Since every polarization of $A' \otimes \overline{\mathbb{F}_p}$ is defined over $\mathbb{F}_{p^2}$ and the set Σ_c is nonempty, there exists a polarization λ_1 of A_1 such that $\ker \lambda_1 = A_1[F]$. Take

$$(3.1) \quad (A', \lambda') := \text{Res}_{\mathbb{F}_{p^2}/\mathbb{F}_p}(A_1, \lambda_1).$$

Then (A', λ') is a superspecial polarized abelian variety over $\mathbb{F}_p$ of dimension n such that

$$(3.2) \quad (A', \lambda') \otimes \mathbb{F}_{p^2} = (A_1, \lambda_1) \times (A_1^{(p)}, \lambda_1^{(p)}).$$

This gives

$$(3.3) \quad \ker \lambda' = \ker \lambda_1 \times \ker \lambda_1^{(p)} = A_1[F] \times A_1^{(p)}[F] = A'[F].$$

From (3.2) we get $\pi_{A'}^2 = -p$. ■

Lemma 3.2. *Assume that $\left(\frac{-1}{p}\right) = 1$. Then for any even positive integer n , there exists a superspecial polarized abelian variety (A, λ) in $\tilde{\Sigma}'_n(\mathbb{F}_p)$*

PROOF. It suffices to prove this for $n = 2$ as we can take the product of copies of such a polarized abelian surface. Let $(A_0, \lambda_0) = (E_0^2, \lambda_0)$ be the superspecial principally polarized abelian surface over $\mathbb{F}_p$ as in Subsection 2.1; one has $\pi_{A_0}^2 = -p$. Consider isogenies $\alpha : (A, \lambda) \rightarrow (A_0, \lambda_0)$ of degree p with $\alpha^* \lambda_0 = \lambda$. The

family $\{\alpha\}$ forms a projective space $\mathbf{P}^1$ that has an $\mathbb{F}_p$ -structure induced from the $\mathbb{F}_p$ -structure of A_0 . If an isogeny $\alpha : (A, \lambda) \rightarrow (A_0, \lambda_0)$ corresponds to a point $[a : b] \in \mathbf{P}^1(\overline{\mathbb{F}}_p)$, then $(A, \lambda) \in \Sigma_2$ if and only if $a^{p+1} + b^{p+1} = 0$ ([4, (3.4), p. 119] and [6, Lemma 4.3]). Since $(-1/p) = 1$, there exists a point $[a : b] \in \mathbf{P}^1(\mathbb{F}_p)$ such that $a^{p+1} + b^{p+1} = a^2 + b^2 = 0$. Then the corresponding isogeny $\alpha : (A, \lambda) \rightarrow (A_0, \lambda_0)$ is defined over $\mathbb{F}_p$ and satisfies both $\pi_A^2 = -p$ and $\ker \lambda = A[F]$. ■

Corollary 3.3. *Assume that $4|n$ or $\left(\frac{-1}{p}\right) = 1$. Then the set $\tilde{\Sigma}'_n(\mathbb{F}_p)$ is nonempty. Consequently, so is the set $\tilde{\Sigma}_n(\mathbb{F}_p)$.*

Remark 3.4. We expect that the sufficient conditions in Corollary 3.3 are also necessary for non-emptiness of $\tilde{\Sigma}'_n(\mathbb{F}_p)$.

For the rest of this section we construct a polarized abelian surface as in Theorem 1.2.

Step 1. There exists a principally polarized abelian surface (A_0, λ_0) over $\mathbb{F}_p$ with $\pi_{A_0}^2 = p$.

Choose an supersingular elliptic curve E over $\mathbb{F}_{p^2}$ such that $\pi_E = p$. Take

$$A_0 = \text{Res}_{\mathbb{F}_{p^2}/\mathbb{F}_p} E.$$

Then the Frobenius endomorphism π_{A_0} of A_0 satisfies $\pi_{A_0}^2 = p$, and one has $A_0 \otimes \mathbb{F}_{p^2} = E \times {}^{\sigma_p}E$. As E admits a principal polarization μ , the product $\lambda_0 := \mu \times {}^{\sigma_p}\mu$ on $A_0 \otimes \mathbb{F}_{p^2}$ is a principal polarization that is defined over $\mathbb{F}_p$.

Step 2. The (covariant) Dieudonné module M_0 of A_0 is a free module over R of rank 2, where $R := \mathbb{Z}_p[\pi_{A_0}] = \mathbb{Z}_p[\sqrt{p}]$. The Frobenius map F and Verschiebung V operate as the multiplication by $\sqrt{p}$. The quasi-polarization induced by λ_0 is a perfect alternating pairing $\langle \cdot, \cdot \rangle : M_0 \times M_0 \rightarrow \mathbb{Z}_p$ such that $\langle ax, y \rangle = \langle x, ay \rangle$ for all $a \in R$ and $x, y \in M_0$. We claim that there exists an R -basis e_1, e_2 for M_0 such that

$$(3.4) \quad \langle e_1, e_2 \rangle = \langle \sqrt{p}e_1, \sqrt{p}e_2 \rangle = 0 \quad \text{and} \quad \langle e_1, \sqrt{p}e_2 \rangle = \langle \sqrt{p}e_1, e_2 \rangle = 1.$$

That is, $\{e_1, \sqrt{p}e_2, \sqrt{p}e_1, e_2\}$ is a Lagrangian $\mathbb{Z}_p$ -basis for the symplectic pairing $\langle \cdot, \cdot \rangle$ on M_0 .

Let $K := \mathbb{Q}_p[\sqrt{p}]$ be the fraction field of R . Let

$$\langle \cdot, \cdot \rangle_K : M_0 \times M_0 \rightarrow R^\vee, \quad R^\vee := \{x \in K \mid \text{tr}_{K/\mathbb{Q}_p}(xR) \subset \mathbb{Z}_p\} = (2\sqrt{p})^{-1}R.$$

be the unique R -bilinear alternating form such that $\langle x, y \rangle = \text{tr}_{K/\mathbb{Q}_p} \langle x, y \rangle_K$, where $\text{tr}_{K/\mathbb{Q}_p}$ denotes the reduced trace from K to $\mathbb{Q}_p$. Put $\psi_K(x, y) := (2\sqrt{p}) \langle x, y \rangle_K$. Then $\psi_K : M_0 \times M_0 \rightarrow R$ is a perfect alternating R -bilinear pairing. Choose an R -basis e_1, e_2 for M_0 such that $\psi_K(e_1, e_2) = 1$. Using the formula $\langle x, y \rangle = \text{tr}_{K/\mathbb{Q}_p} (2\sqrt{p})^{-1} \psi_K(x, y)$, we check (3.4) as follows:

$$\begin{aligned} \langle \sqrt{p}e_1, e_2 \rangle &= \langle e_1, \sqrt{p}e_2 \rangle = \text{tr}(2\sqrt{p})^{-1} \sqrt{p} = \text{tr } 2^{-1} = 1, \\ \langle e_1, e_2 \rangle &= \text{tr}(2\sqrt{p})^{-1} = 0, \quad \langle \sqrt{p}e_1, \sqrt{p}e_2 \rangle = \langle e_1, pe_2 \rangle = 0. \end{aligned}$$

Step 3. Let $\alpha : A \rightarrow A_0$ be any isogeny of degree p defined over $\mathbb{F}_p$, and let $\lambda := \alpha^* \lambda_0$. Then A is a superspecial abelian surface over $\mathbb{F}_p$ with $\pi_A^2 = p$, and λ has the property $\ker \lambda = A[F]$.

The Dieudonné module M of A fits into $\mathbf{V}M_0 = \sqrt{p}M_0 \subset M \subset M_0$ with $\dim M/M_0 = 1$. Clearly, M is an R -submodule so that F^2 acts as p on M , and M is superspecial. Put $\overline{M}_0 := M_0/\mathbf{V}M_0 = \text{Span}\{\bar{e}_1, \bar{e}_2\}_{\mathbb{F}_p}$. Define a pairing

$$(\cdot, \cdot) : M_0 \times M_0 \rightarrow \mathbb{Z}_p, \quad (x, y) := \langle x, Fy \rangle.$$

Clearly, $(M_0, \mathbf{V}M_0) = (\mathbf{V}M_0, M_0) = p\mathbb{Z}_p$. Modulo p , one obtains an $\mathbb{F}_p$ -bilinear pairing $(\cdot, \cdot) : \overline{M}_0 \times \overline{M}_0 \rightarrow \mathbb{F}_p$ satisfying $(\bar{e}_1, \bar{e}_2) = -(\bar{e}_2, \bar{e}_1) = 1$ and $(\bar{e}_1, \bar{e}_1) = (\bar{e}_2, \bar{e}_2) = 0$. Now it is not hard to see that the condition $\ker \lambda = A[F]$ is equivalent to $\mathbf{V}(M^t) = M$, which is also equivalent to the condition $(\overline{M}, \overline{M}) = 0$ (see the proof of [6, Lemma 4.3]). As $\overline{M}$ is generated by a vector $v = a\bar{e}_1 + b\bar{e}_2$, where $a, b \in \mathbb{F}_p$, the condition $(\overline{M}, \overline{M}) = 0$ follows immediately from $(v, v) = 0$. Thus, one proves $\ker \lambda = A[F]$.

This completes the proof of Theorem 1.2

ACKNOWLEDGEMENTS

The author is grateful to Tomoyoshi Ibukiyama for kind explanations of his recent works [1] and [2]. The paper is completed during the author's visit at Münster University; he acknowledges Urs Hartl and the institution for warm hospitality and excellent working conditions. The author is partially supported by the grant 104-2115-M-001-001MY3.

REFERENCES

- [1] T. Ibukiyama, Quinary lattices and binary quaternion hermitian lattices. Preprint, 2016, 14 pp.
- [2] T. Ibukiyama, Type numbers of quaternion hermitian forms and supersingular abelian varieties. Preprint, 2016, 18 pp.
- [3] T. Ibukiyama and T. Katsura, On the field of definition of superspecial polarized abelian varieties and type numbers. *Compositio Math.* **91** (1994), 37–46.
- [4] T. Katsura and F. Oort, Families of supersingular abelian surfaces, *Compositio Math.* **62** (1987), 107–167.
- [5] K.-Z. Li and F. Oort, *Moduli of Supersingular abelian Varieties*. Lecture Notes in Math., vol. 1680, Springer-Verlag, 1998.
- [6] C.-F. Yu, The supersingular loci and mass formulas on Siegel modular varieties. *Doc. Math.* **11** (2006), 449–468.
- [7] C.-F. Yu, An explicit reciprocity law arising from superspecial abelian varieties and trace formulas. arXiv:1210.1120, 40 pp.

INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, 6TH FLOOR, ASTRONOMY MATHEMATICS BUILDING, NO. 1, ROOSEVELT RD. SEC. 4, TAIPEI, TAIWAN, 10617

E-mail address: chiafu@math.sinica.edu.tw

NATIONAL CENTER FOR THEORETICAL SCIENCES NO. 1 ROOSEVELT RD. SEC. 4, NATIONAL TAIWAN UNIVERSITY TAIPEI, TAIWAN, 10617